

1 Sums and Limits

mathclap & friends

$$X = \sum_{1 \leq i \leq j \leq n} X_{ij}$$

$$X = \sum_{1 \leq i \leq j \leq n} X_{ij}$$

$$X = \sum_{1 \leq i \leq j \leq n} X_{ij}$$

$$X = \sum_{1 \leq i \leq j \leq n} X_{ij}$$

Cramped

$$x^2 \leftrightarrow x^2 \quad x^2 \leftrightarrow x^2$$

Smashoperator

$$V = \sum_{1 \leq i \leq j \leq n}^{\infty} V_{ij} \quad X = \sum_{1 \leq i \leq j \leq n}^{3456} X_{ij} \quad Y = \sum_{1 \leq i \leq j \leq n} Y_{ij} \quad Z = \sum_{1 \leq i \leq j \leq n} T Z_{ij}$$

$$V = \sum_{1 \leq i \leq j \leq n}^{\infty} V_{ij} \quad X = \sum_{1 \leq i \leq j \leq n}^{3456} X_{ij} \quad Y = \sum_{1 \leq i \leq j \leq n} Y_{ij} \quad Z = \sum_{1 \leq i \leq j \leq n} T Z_{ij}$$

Adjustlimits

a) $\lim_{n \rightarrow \infty} \max_{p \geq n}$ b) $\lim_{n \rightarrow \infty} \max_{p^2 \geq n}$ c) $\lim_{n \rightarrow \infty} \sup_{p^2 \geq nK}$ d) $\limsup_{n \rightarrow \infty} \max_{p \geq n}$

a) $\lim_{n \rightarrow \infty} \max_{p \geq n}$ b) $\lim_{n \rightarrow \infty} \max_{p^2 \geq n}$ c) $\lim_{n \rightarrow \infty} \sup_{p^2 \geq nK}$ d) $\limsup_{n \rightarrow \infty} \max_{p \geq n}$

2 Tags

$$a = b$$

Q&A

See Q&A or is it better with Q&A?

$$a = b$$

Q&A

$$a = b$$

[Q&A]

Normal tags.

$$a = a$$

(1)

That was equation (1).

OK tags.

$$a = a \quad [2]$$

That was equation [2], but recall [1]

odd tag.

$$a = a \quad \{3\}$$

That was equation {3}, but recall {1} and {2}.

weird tag.

$$b = b \quad ((4))$$

That was equation ((4)), but recall ((1)), ((2)) and ((3)).

Normal tags again.

$$c = c \quad (5)$$

Non-textual

$$d = d \quad (n^{th})$$

That was equation (5), but recall (1), (2), (3), (4) and (nth).

$$\begin{aligned} a &= a \\ b &= b \end{aligned} \quad (6) \quad (**)$$

This should refer to the equation containing $a = a$: (6). Then a switch of tag forms.

$$c = c \quad (7)$$

$$d = d \quad (8)$$

This should refer to the equation containing $d = d$: (8) (but recall (6)).

$$e = e \quad (9)$$

$$f = f \quad (10)$$

$$1 + 1 = 2$$

$$2 + 2 = 4$$

Blabla (2).

3 Arrows

$$\begin{array}{c}
A \xrightleftharpoons[\text{under}]{\text{over}} B \xrightleftharpoons[\text{under}]{\text{over}} C \\
x \xrightleftharpoons[\text{under}]{\text{overlooooooong}} y \xrightleftharpoons[\text{underlooooooong}]{\text{over}} z \\
x \xrightleftharpoons[\substack{\text{bar} \\ \text{bluuuuuuuuub}}]{\substack{\text{foo} \\ \text{baz}}} y \xrightleftharpoons[\substack{\text{blaaaaaab} \\ \text{heeeeee}}]{\text{baz}} t \xrightarrow{\text{heeeeee}} k \\
k \xleftarrow{\substack{\text{.} \\ \text{...}}} l \xleftarrow{\substack{\text{.} \\ \text{...}}} m \xrightarrow{\substack{\text{.} \\ \text{...}}} n \xrightarrow{\substack{\text{.} \\ \text{...}}} o \\
x \xrightleftharpoons[\substack{\text{blaaaaaab} \\ \text{bla}}]{\text{bluuuuub}} y \xrightleftharpoons[\substack{\text{bla} \\ \text{aa}}]{\text{bla}} z \\
z = \underbrace{x}_{\substack{\text{real} \\ \text{.}}} + i \underbrace{y}_{\substack{\text{imaginary} \\ \text{.}}} \underbrace{1+1}_{=2} \text{ complex number}
\end{array}$$

4 Matrices

$$\begin{array}{c}
c \quad cocococococo \\
c \quad \quad c \\
lalalalalala \quad l \\
l \quad \quad l \\
rererererere \quad r \\
r \quad r \\
\begin{pmatrix} ppppppp & foo \\ l & ppppppppppppppp \end{pmatrix} \\
\begin{bmatrix} b & b \\ b & b \end{bmatrix} \\
\begin{Bmatrix} B & B \\ B & BBBBBrBBBBB \end{Bmatrix} \\
\begin{vmatrix} v & v \\ v & v \end{vmatrix} \\
\begin{Vmatrix} V & V \\ VVVVVVcVVVVVV & bar \end{Vmatrix} \\
\left| \begin{smallmatrix} a & blblblblbbblblblbl \\ c & d \end{smallmatrix} \right| \\
\begin{bmatrix} a & -b \\ -c & d \end{bmatrix} \begin{bmatrix} a & -b \\ -c & d \end{bmatrix}
\end{array}$$

$$\begin{aligned}
& \left\| \begin{pmatrix} e & -f \\ -g & h \end{pmatrix} \right\| \left\| \begin{pmatrix} e & -f \\ -g & h \end{pmatrix} \right\| \\
& \left[\begin{pmatrix} a & -bbbb \\ -c & d \end{pmatrix} \right] \left[\begin{pmatrix} a & -bbbb \\ -c & d \end{pmatrix} \right] \\
& \left\| \begin{pmatrix} e & -fffff \\ -g & h \end{pmatrix} \right\| \left\| \begin{pmatrix} e & -fffff \\ -g & h \end{pmatrix} \right\| \\
& \left[\begin{pmatrix} a & -bbbb \\ -c & d \end{pmatrix} \right] \left[\begin{pmatrix} a & -bbbb \\ -c & d \end{pmatrix} \right] \\
& \left\| \begin{pmatrix} e & -fffff \\ -g & h \end{pmatrix} \right\| \left\| \begin{pmatrix} e & -fffff \\ -g & h \end{pmatrix} \right\|
\end{aligned}$$

5 Cases

$$\begin{aligned}
& \begin{cases} E = mc^2 & \text{Nothing to see here} \\ \int x - 3 dx & \text{Integral is text style} \end{cases} \\
& \begin{cases} E = mc^2 & c \approx 3.00 \times 10^8 \text{ m/s} \\ \int x - 3 dx & \text{Integral is display style} \end{cases} \\
& a = \begin{cases} E = mc^2 & \text{Nothing to see here (text in math)} \\ \int x - 3 dx & \text{Integral is display style (text in math)} \end{cases} \\
& \begin{cases} E = mc^2 & 5^6 \text{ and so on} \\ \int x - 3 dx & \int x dx \end{cases} = b \\
& \begin{cases} x^2 & \text{for } \int x dx > 0 \\ x^3 & \text{else} \end{cases} \Rightarrow \dots \\
& \begin{cases} E = mc^2 & 5^6 \text{ and so on} \\ \int x - 3 dx & \int x dx \end{cases} = b \\
& \begin{cases} x^2 & \text{for } \int x dx > 0 \\ \int x^3 x & \text{else} \end{cases} \Rightarrow \dots \\
& \text{foo} = \begin{cases} \pi & \text{if something} \\ \int \Omega^\Xi \Omega & \text{otherwise} \end{cases}
\end{aligned}$$

6 Gathered

$$A = \boxed{\text{first}} B$$
$$\boxed{\text{last}}$$

$$a = b + c$$

$$b = c + d$$

...

$$\boxed{\text{hello}}$$

$$\boxed{f(x) = \int h(x) dx}$$

$$= g(x)$$

$$a = b$$

(11)

Some text

$$c = d$$

(12)

Some short text

$$e = f$$

(13)

7 Delimiters

$$\begin{array}{|c|c|c|} \hline \left| \frac{a}{c} \right| & \left| \frac{a}{c} \right| & \left| \frac{a}{b} \right| \\ \hline \left| \frac{a}{b} \right| & \left| \frac{a}{b} \right| & \left| \frac{a}{b} \right| \\ \hline \end{array}$$
$$|\pi| \quad |\neg\phi|$$

$$\left\langle A, \frac{1}{2} \right\rangle \quad \left\langle B \middle| \sum_k f_k \middle| C \right\rangle$$

$$\left\{ x \in X \middle| \frac{\sqrt{x}}{x^2 + 1} > 1 \right\}$$

$$\langle 1 \mid \frac{8}{4} \mid 3 \rangle \quad \left\langle 1 \middle| \frac{8}{4} \middle| 3 \right\rangle \quad \langle 1 \mid \frac{8}{4} \mid 3 \rangle$$

$$\left(\frac{\pi}{\omega} \right) \cdot \left[\int x dx \right] \dots [\sqrt{\frac{\sin x}{\cos z}}] \dots (\frac{\frac{foo}{bar}}{\frac{baz}{qux}})$$

Operators

$$a := b \quad a := b \quad a := b$$

$$a := b \quad c ::\approx d \quad e :: f$$

$$\times \times \dagger \otimes \otimes$$

8 Prescripts

$${}^4_{12}\mathbf{C}_2^{5+} \quad {}^{14}{}_2\mathbf{C}_2^{5+} \quad {}^4_{12}\mathbf{C}_2^{5+} \quad {}^{14}{}_2\mathbf{C}_2^{5+} \quad {}_2\mathbf{C}_2^{5+}$$

$${}^A_Z\mathbf{X} \rightarrow {}^{A-4}_{Z-2}\mathbf{Y} + {}^4_2\alpha$$

$$a = \frac{xy + xy + \int xy \, dx + xy + xy}{z} = \frac{xy + xy + \int xy \, dx + xy + xy}{z}$$

9 Multlines

$$\begin{aligned} p(x) = & 3x^6 + 14x^5y + 590x^4y^2 + 19x^3y^3 \\ & - 12x^2y^4 - 12xy^5 + 2y^6 - a^3b^3 \end{aligned}$$

$$A = \boxed{\begin{matrix} first \\ \dots \\ last \end{matrix}} \quad B$$

$$A = \boxed{\begin{matrix} first \\ \dots \\ last \end{matrix}} \quad B$$

$$A = \quad \boxed{\begin{matrix} first \\ \dots \\ last \end{matrix}} \quad B$$

$$A = \boxed{\begin{matrix} first \\ \dots \\ last \end{matrix}} \quad B$$

$$A = \boxed{\begin{matrix} first \\ \dots \\ last \end{matrix}} \quad B$$

$$A = \boxed{first}$$

$$A = \boxed{first} \qquad \qquad B \\ \boxed{last}$$

$$A = \qquad \qquad \boxed{last}B$$

$$A = \boxed{first} \qquad B \\ \boxed{last}$$

$$foo ::= x = 1, \quad x + 1 = 2 \quad (14) \\ y = 2$$

$$x = 1, \quad x + 1 = 2 \\ bar ::= \qquad \qquad \qquad y = 2 \quad (15)$$

10 Spread-lines

Spread it

$$\begin{array}{ccc} a & b & c \\ d & e & f \\ g & h & i \end{array}$$

$$\begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,n} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1} & a_{m,2} & \cdots & a_{m,n} \end{pmatrix}$$

$$\begin{cases} \frac{a}{c} \frac{b}{d} \\ n/2 & \text{if } n \text{ is even} \\ -(n+1)/2 & \text{if } n \text{ is odd} \end{cases}$$

$$a=b+c-d$$

$$+e-f$$

$$\qquad\qquad\qquad(16)$$

$$=g+h$$

$$= i$$

$$a+b+c+d+e+f$$

$$+i+j+k+l+m+n \quad (17)$$

$$a=b \qquad\qquad\qquad(18)$$

$$c=d \qquad\qquad\qquad(19)$$

$$a_1=b_1+c_1 \qquad\qquad\qquad(20)$$

$$a_2=b_2+c_2-d_2+e_2 \qquad\qquad\qquad(21)$$

$$a_{11}=b_{11}$$

$$a_{12}=b_{12}$$

$$a_{21}=b_{21}$$

$$a_{22}=b_{22}+c_{22}$$

$$8\\$$

$$x = y_1 - y_2 + y_3 - y_5 + y_8 - \dots \quad \text{by foo} \quad (22)$$

$$= y' \circ y^* \quad \text{by baz} \quad (23)$$

$$= y(0)y' \quad \text{by Axiom 1.} \quad (24)$$

$$\left. \begin{array}{l} B' = -\partial \times E, \\ E' = \partial \times B - 4\pi j, \end{array} \right\} \text{Maxwell's equations}$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$\sum_{\substack{i \in \Lambda \\ 0 < j < n}} P(i, j)$$

$$y = ax^2 + bx + c \quad (25)$$

$$f(x) = x^2 + 2xy + y^2 \quad (26)$$

<div style="border: 1px solid black; padding: 2px; display: inline-block;">Firstline</div>	<div style="border: 1px solid black; padding: 2px; display: inline-block;">Secondline</div>
$L + E + F + T$	$R + I + G + H + T$
$L + E + F + T$	$R + I + G + H + T$
<div style="border: 1px solid black; padding: 2px; display: inline-block;">WupWup</div>	<div style="border: 1px solid black; padding: 2px; display: inline-block;">Lastline</div>

11 Stepped lines

1* $x = 1, \quad x + 1 = 2$ over
2* $y = 2$ over

42

See: $s = 2.8, \quad s + 0.2 = 3$ the end
See: $t = 4.5$ the end

1337

12 Shifting equations

Part 1
$=$ [] 2nd line
$19 +$ [] last part

$$\Downarrow \quad [1] = [2] \quad (27)$$

$$\Downarrow \quad [3] = [4] \quad (28)$$

$$\begin{aligned} a &= b \\ &\vdots \\ &= c \\ &\vdots \\ &= d \end{aligned}$$